

PROVEN COCCIDIOSTAT IN NEW FORMULATION

COXIDIN® (monensin sodium)

Coxidin® 200 MicroGranulate

- A coccidiostatic additive for broilers and turkeys
- Contains 200 gm of monensin sodium per kg in Coxidin.
- It is available as homogenous granules.

EFFICACY

- Efficacious in the early stage of the coccidia lifecycle (schizogony) which prevents damage to the epithelial cells of the gut lining.
- Helps build up natural immunity.
- Significantly reduces the risk of secondary infection by pathogens and minimizes weight loss and poor feed conversion.
- Coxidin® 200 is a broad spectrum coccidiostat.
- Coxidin® 200 is a Polyether ionophore.



Coxidin® 200

- Forms complex with monovalent cations & shows swelling
- Creates huge intrasporozoites vacuoles
- Causes extreme osmotic damage
- Finally kills off the sporozoite

SAFE

- Very well tolerated.
- Does not adversely affect feed and water intake.

FREE-FLOWING

- Unique granulated formulation.
- Excellent mixability and homogeneity.
- Remains stable during pelletisation.

ACTIVITY

Coxidin® 200 is effective against the main coccidia species:

- *Eimeria tenella*, *E. acervulina*, *E. mivati*, *E. brunetti*, *E. maxima*, *E. necatrix* in chicken.
- *E. adenoides*, *E. meleagrimitis* and *E. gallopavonis* in turkeys.

DOSAGE

SPECIES	Coxidin® 200 Premix gm/ton Feed	Content of monensin complete feedingstuff (ppm)	Duration of Feeding
Chicken For Fattening	500-625	100-125	Entire rearing period
Chicken For Laying	500-625	100-125	Up to 16 weeks of age
Turkeys	300-500	60-100	Up to 16 weeks of age

WITHDRAWAL PERIOD

Chicken & turkeys: 1 day after the last administration of the preparation.

INTERACTIONS

Coxidin® 200 microgranulate is compatible with most feed additives: antibiotics, vitamins and minerals. It is not compatible with tiamulin, erythromycin, oleandomycin, sulphadimethoxine, sulpha- dimidine and sulphaquinoxaline.

STORAGE

Store in the original package in a closed, dry and well-ventilated area at room temperature, protected from direct sunlight.

PACKAGING

25 kg PE-Lined multi-layered paper bags.

Huvepharma AD
3a Nikolay Haytov Str., 1113 Sofia, Bulgaria
P +359 2 862 5331 • F +359 2 862 5334
sales@huvepharma.com



Coxidin® is Registered trademark of Huvepharma

© 2000 Huvepharma AD

Huvepharma N.V.
Uitbreidingstraat 80 • 2600 Antwerp • Belgium
P +32 3 288 1849 • F +32 3 289 7845
customerservice@huvepharma.com