


ANIMAL FEED SUPPLEMENT



# COYDEN<sup>®</sup> 12.5%

clopidol

AUTHENTIC  
EUROPEAN  
SOURCE



## Coyden® 12.5%- Highly effective in the prevention of coccidiosis

Coyden® 12.5% is a non-dusting, granular premix which contains 125mg/g of the active ingredient clopidol (formulated from meticlorpindol). It was first marketed in 1968 and was proven safe and effective for more than 2 decades in the field of coccidiosis prevention.

Today, poultry producers are pleading for additional coccidiosis solutions, which is one of the reasons for the revival of historically successful products to be applied in rotation programs.

### BENEFITS

- Proven **broad spectrum** efficacy
- **Unique** chemistry limits possibility of cross resistance to other anticoccidials
- **No known incompatibilities** with other feed additives and medications (like tiamulin hydrogen fumarate)
- **ideal for clean-up** low lesion scores and low oocyst excretion



### DOSAGE

Target species	Minimum and maximum content of clopidol in complete feedingstuff (ppm)	Quantity of Coyden® 12.5% incorporated into the feedingstuff (g/ton)	Maximum age	Withdrawal period before slaughter
Chickens for fattening	125	1000	-	5 days
Breeders & Layers	125	1000	14 weeks	5 days

