

THE NEW COMBINATION COCCIDIOSTAT FOR BROILER CHICKENS AND TURKEYS

Monimax[®]
(monensin/nicarbazin)


Pharmacological action

Monimax[®] is a coccidiostat containing the active ingredients monensin and nicarbazin. The combined strength of the ionophore monensin and the chemical nicarbazin leads to a unique new product in coccidiosis control.

Monimax[®] is effective against coccidiosis caused by all major *Eimeria species* in broilers and turkeys.

On cellular level the effect of Monimax[®] to inhibit the development of sporozoites and trophozoites is enhanced by the synergetic action of monensin and nicarbazin.





Composition

| Content | Monimax® |
|---------------------------------------|---------------|
| Monensin sodium | 8 g per 100 g |
| Nicarbazin | 8 g per 100 g |
| Starch, wheat bran, calcium carbonate | Up to 100 g |

Indications

As an aid in the prevention of coccidiosis in broiler chickens, rearing pullets and turkeys caused by *Eimeria tenella*, *E. acervulina*, *E. maxima*, *E. necatrix*, *E. brunetti*, *E. mitis* and *E. praecox* in chickens for fattening and *E. meleagrimitis*, *E. adenoides*, *E. gallopavonis*, *E. meleagridis* and *E. dispersa* in turkeys.

Contraindications

Do not feed to laying hens. Do not mix with other anticoccidials.

Target species

Broiler chickens, rearing pullets and turkeys.

Mode of administration

Orally, well homogenized into the feed. Monimax® should be thoroughly and evenly mixed in the feed in accordance with current good manufacturing practice for medicated feed.

Dosage

For prevention of coccidiosis in broilers and turkeys. Monimax® is most frequently used at a concentration of **40/40 ppm**. This can be achieved by blending **500 g** of Monimax® per ton of feed. In case of severe infection the dosage can be increased up to **50/50 ppm** or **625 g** Monimax® per ton of feed.

Instructions for use

Monimax® microgranules must be mixed with other ingredients before feeding in order to have an optimal homogeneity and to avoid segregation. Mixing and conveying equipment should be properly cleaned to prevent carry-over.

Side effects

None under correct use of the product.

Withdrawal period

Check with your local Huvepharma representative.

Storage

In the original packing, well closed, in dry and well-ventilated facilities, protected from direct sunlight.

Packaging

Monimax® is packed in 20 kg bags.

Warning

Dangerous for equines.

Poultry should not be treated with tiamulin for at least seven days before or after receiving feed containing Monimax®.

When mixing, direct contact with the skin and mucous membranes should be avoided. Accidental ingestion should be avoided. Wear overalls, safety glasses, mask and impervious gloves when handling or mixing the product. Contaminated clothing should be removed and any splashes on to the skin should be washed off immediately. If accidental eye contact occurs, immediately rinse thoroughly with water. Seek medical advice if irritation persists. Wash hands after use.

| | | broiler chickens | rearing pullets | turkeys |
|---|---------|------------------|-----------------|---------|
| Content of monensin/nicarbazin in complete feedingstuff (ppm) | minimum | 40/40 | 40/40 | 40/40 |
| | maximum | 50/50 | 50/50 | 50/50 |
| Quantity of Monimax® incorporated in feed (g per ton) | minimum | 500 | 500 | 500 |
| | maximum | 625 | 625 | 625 |

Huvepharma EAD

3^a Nikolay Haytov Str., 1113 Sofia, Bulgaria
P +359 2 862 5331 • F +359 2 862 5334
sales@huvepharma.com



© 2015 Huvepharma EAD. All rights reserved.
Monimax.0615.EN003/Novis

Huvepharma N.V.

Uitbreidingstraat 80 • 2600 Antwerp • Belgium
P +32 3 288 1849 • F +32 3 289 7845
customerservice@huvepharma.com