

THE NEW COMBINATION COCCIDIOSTAT FOR BROILER CHICKENS

PHARMACOLOGICAL ACTION

Nimax® is a microGranulated coccidiostat containing 2 active ingredients. The combined strength of the chemical **nicarbazin** and the ionophore **maduramicin** leads to a unique new product in coccidiosis control.

Nimax® is effective against coccidiosis caused by all major Eimeria species in broilers. On cellular level the effect of Nimax® to inhibit the development of sporozoites and trophozoites is enhanced by the **synergetic** action of nicarbazin and maduramicin.

This innovative combination product will not only result in **superior coccidiosis control**, but will also **enhance performance levels**.

Nimax[®]
(nicarbazin/maduramicin)

- nicarbazin: 80 gm / kg
- maduramicin: 7.5 gm/kg



HUVEPHARMA[®]

We add performance to your business



COMPOSITION

Content	NIMAX®
Nicarbazin	8 gm per 100 gm
Maduramicin	0.75 gm per 100 gm

INDICATIONS

As an aid in the prevention of coccidiosis caused by *Eimeria tenella*, *E.acervulina*, *E. maxima*, *E. necatrix*, *E. brunetti*, *E. mitis* and *E. praecox* in chickens.

CONTRAINDICATIONS

Do not feed to horses or other equines. Do not mix with other coccidiostats.

TARGET SPECIES

Broiler Chickens.

MODE OF ADMINISTRATION

Orally, well homogenized into the feed. Nimax® should be thoroughly and evenly mixed in the feed in accordance with current good manufacturing practice for medicated feed.

DOSAGES

For prevention of coccidiosis in broilers.

Nimax® is used at an inclusion rate of **500 gm per ton of feed**

		Broiler chickens
Content of nicarbazin/ maduramicin in complete feedingstuff (ppm)	maximum	40/3.75
Quantity of Nimax® incorporated in feed (gm per ton)	maximum	500

INSTRUCTIONS FOR USE

Nimax® microgranules must be mixed with other ingredients before feeding in order to have an optimal homogeneity and to avoid segregation. Mixing and conveying equipment should be properly cleaned to prevent carry-over.

SIDE EFFECTS

None under correct use of the product.

WITHDRAWAL PERIOD

None under correct use of the product.

STORAGE

In the original packing, well closed, in dry and well ventilated facilities, protected from direct sunlight.

PACKING

Nimax® is packed in 25 kg bags.

WARNING

Dangerous for equines.

When mixing, direct contact with the skin and mucous membranes should be avoided. Accidental ingestion should be avoided. Wear overalls, safety glasses, mask and impervious gloves when handling or mixing the product. Contaminated clothing should be removed and any splashes on to the skin should be washed off immediately. If accidental eye contact occurs, immediately rinse thoroughly with water. Seek medical advice if irritation persists. Wash hands after use.

Huvepharma AD

3a Nikolay Haytov Str., 1113 Sofia, Bulgaria

P +359 2 862 5331 . F +359 2 862 5334

sales@huvepharma.com

