

ANIMAL FEED SUPPLEMENT



Poulcox[®] 15%

Monensin sodium



HUVEPHARMA[®]

Poulcox® 15% - Prevention of coccidia in chicken

POULCOX 150 MICROGRANULATE

- A coccidiostatic feed additive for chicken.
- Contains 150 gm of monensin sodium per kg in Poulcox.
- It is available as homogenous granules.

BENEFITS

- Positive effect on **litter quality**
- **Prevention of** *E. acervulina*, *E. maxima* & *E. tenella* in adverse environmental conditions.
- **Ease of use:** can be used in chickens for fattening, chickens reared for laying & breeding stock.



DOSAGE

Target species	Minimum and maximum dose of monensin (ppm)	Recommended dose of Poulcox® 15% (g/ton of feed)	Maximum age	Withdrawal period before slaughter
Chickens for fattening	100 - 125	665 - 835	Entire rearing period	1 day
Chickens reared for laying & breeding stock	100 - 125	665 - 835	18 weeks	1 day

WITHDRAWAL PERIOD

Chicken : 1 day after the last administration of the preparation.

STORAGE

Store in the original package in a closed, dry and well-ventilated area at room temperature, protected from direct sunlight.

SHELF LIFE

2 Years.

PACKAGING

25 kg PE-Lined multi-layered paper bags.